

Anthropometric and dietary indicators applied in population-based surveys: a systematic review

Indicadores antropométricos e dietéticos utilizados em estudos de base populacional: uma revisão sistemática

Indicadores antropométricos y dietéticos utilizados en estudios poblacionales: una revisión sistemática

Received: 01/06/2022 | Reviewed: 01/16/2022 | Accept: 01/23/2022 | Published: 01/25/2022

Natália Louise de Araújo Cabral

ORCID: <https://orcid.org/0000-0002-9166-1917>

Federal Institute of Education, Science and Technology of Sertão Pernambucano, Brazil

E-mail: natalia.louise@ifsertao-pe.edu.br

Nila Patrícia Freire Pequeno

ORCID: <https://orcid.org/0000-0003-1279-2554>

Federal University of Rio Grande do Norte, Brazil

E-mail: nilapfp@hotmail.com

David Franciole de Oliveira Silva

ORCID: <https://orcid.org/0000-0003-0940-1356>

Federal University of Rio Grande do Norte, Brazil

E-mail: davfranci@hotmail.com

Sandra Patrícia Crispim

ORCID: <https://orcid.org/0000-0002-2257-9899>

Federal University of Paraná, Brazil

E-mail: sandracrispim@gmail.com

Dirce Maria Lobo Marchioni

ORCID: <https://orcid.org/0000-0002-6810-5779>

University of São Paulo, Brazil

E-mail: marchioni@usp.br

Severina Carla Vieira Cunha Lima

ORCID: <https://orcid.org/0000-0001-8268-1986>

Federal University of Rio Grande do Norte, Brazil

E-mail: scvclima@gmail.com

Clélia de Oliveira Lyra

ORCID: <https://orcid.org/0000-0002-1474-3812>

Federal University of Rio Grande do Norte, Brazil

E-mail: clelia_lyra@yahoo.com.br

Abstract

Aim: To identify population-based health and nutrition surveys, conducted with adults and the elderly, and performed in the Americas, Europe, and Oceania, to investigate the more common anthropometric and food consumption methods used, their applicability, and their limitations. **Methods:** Electronic databases (LILACS, PubMed, and SCOPUS) were systematically searched for studies published between 1997 and 2017 in Portuguese, English, or Spanish. 48 studies (45.8% carried out in the Americas) met the eligibility criteria and were included in the review. The data were analyzed in 2018. **Results:** The methodological quality of most of the studies (64.4%) was classified as moderate, according to the Agency for Healthcare Research and Quality checklist for cross-sectional studies and the Newcastle-Ottawa scale for cohort studies. 35.4% of the articles evaluated only food consumption, 29.2% just anthropometry, while 35.4% evaluated food consumption and anthropometric measurements. The most used food survey methods were food record (31% of studies) and the 24-hour dietary recall (22% of studies). Body mass index (BMI) was the most used indicator for anthropometric nutritional status assessment. Although most of the studies used the World Health Organization classification criteria, these studies did not adopt the different cut-off points for BMI classification for adults and the elderly. **Conclusion:** BMI and methods that record current consumption, such as the food record and the 24-hour dietary recall, were the main methods of assessing nutritional status, taking into consideration the easy application, low cost, and good reproducibility. Systematic Review Registration: PROSPEROCD42017071392.

Keywords: Nutrition surveys; Population surveys; Food consumption; Anthropometry; Systematic review.

Resumo

Objetivo: Identificar pesquisas de saúde e nutrição de base populacional, com adultos e idosos, e realizadas nas Américas, Europa e Oceania, para investigar os métodos antropométricos e de consumo alimentar mais utilizados, sua aplicabilidade e limitações. **Metodologia:** Bases de dados eletrônicas (LILACS, PubMed e SCOPUS) foram sistematicamente pesquisadas em busca de estudos publicados entre 1997 e 2017 em português, inglês ou espanhol. 48 estudos (45.8% realizados nas Américas) atenderam os critérios de elegibilidade e foram incluídos na revisão. Os dados foram analisados em 2018. **Resultados:** A qualidade metodológica da maioria dos estudos (64,4%) foi classificada como moderada, conforme o checklist da Agency for Healthcare Research and Quality para estudos transversais e a escala Newcastle-Ottawa para estudos de coorte. 35,4% dos artigos avaliaram apenas o consumo alimentar, 29,2% apenas a antropometria, enquanto 35,4% avaliaram o consumo alimentar e as medidas antropométricas. Os métodos de consumo alimentar mais utilizados foram o registro alimentar (31% dos estudos) e o recordatório de 24h (22% dos estudos). O índice de massa corporal (IMC) foi o indicador mais utilizado para avaliação antropométrica. Embora a maioria dos estudos tenha utilizado os critérios da Organização Mundial da Saúde, estes não adotaram pontos de corte diferentes para classificar o IMC de adultos e idosos. **Conclusão:** o IMC e os métodos que registram o consumo atual, como o registro alimentar e o recordatório de 24h, foram os principais métodos de avaliação do estado nutricional, levando-se em consideração a fácil aplicação, baixo custo e boa reprodutibilidade. Registro de revisão sistemática: PROSPEROCD42017071392.

Palavras-chave: Inquéritos nutricionais, Inquéritos populacionais, Consumo alimentar, Antropometria, Revisão sistemática.

Resumen

Objetivo: Identificar encuestas poblacionales de salud y nutrición, realizadas con adultos y ancianos, y realizadas en las Américas, Europa y Oceanía, para investigar los métodos antropométricos y de consumo de alimentos más utilizados, su aplicabilidad y limitaciones. **Metodología:** Se realizaron búsquedas sistemáticas en bases de datos electrónicas (LILACS, PubMed y SCOPUS) de estudios publicados entre 1997 y 2017 en portugués, inglés o español. 48 estudios (45,8% realizados en las Américas) cumplieron con los criterios de elegibilidad y se incluyeron en la revisión. Los datos se analizaron en 2018. **Resultados:** La calidad metodológica de la mayoría de los estudios (64,4%) se clasificó como moderada, según la lista de verificación de la Agency for Healthcare Research and Quality para los estudios transversales y la escala de Newcastle-Ottawa para los estudios de cohortes. 35,4% de los artículos solo evaluó el consumo de alimentos, el 29,2% solo la antropometría, mientras que el 35,4% evaluó el consumo de alimentos y las medidas antropométricas. Los métodos de consumo de alimentos más utilizados fueron el registro de alimentos (31% de los estudios) y el recuerdo de 24 horas (22% de los estudios). El índice de masa corporal (IMC) fue el indicador más utilizado para la evaluación antropométrica del estado nutricional. Aunque la mayoría de los estudios utilizaron los criterios de clasificación de la Organización Mundial de la Salud, estos estudios no adoptaron diferentes puntos de corte para clasificar el IMC de adultos y ancianos. **Conclusión:** El IMC y los métodos que registran el consumo actual, como los registros de alimentos y el retiro de alimentos de 24 horas, fueron los principales métodos para evaluar el estado nutricional, teniendo en cuenta su fácil aplicación, bajo costo y buena reproducibilidad. Registro de revisión sistemática: PROSPEROCD42017071392.

Palabras clave: Encuestas nutricionales; Estudios poblacionales; Consumo alimentario; Antropometría; Revisión sistemática.

1. Introduction

The demographic, epidemiological and nutritional changes, occurring worldwide, in recent decades have resulted in significant transformations in morbidity and mortality patterns. In this context, population-based health and nutrition surveys have been applied since the 1960s in developed countries, and are currently experiencing an increase in many developing countries, with the objective of monitoring the health status of populations (Viacava, 2002).

The health and nutrition surveys are essential for health surveillance, in addition to enabling the analysis of inequalities. These surveys provide updated and reliable information that may contribute to the development of health programs and services in the most diverse population groups, serving as important instruments for public policies planning and evaluation (Barros, 2008; Malta, Célia, & Szwarcwald, 2017; Malta et al, 2008; Sperandio & Priore, 2017).

In most cases, population-based surveys are part of cohort or cross-sectional surveys, and aggregate a set of sociodemographic characteristics, variables of lifestyle factors and health outcomes, data collection methodology, and periodicity. A periodic update on survey is important for the consolidation of the information collected, which becomes the

basis for population references, enabling surveillance of chronic noncommunicable diseases (NCDs) and their causes (Szwarcwald et al., 2014).

Health-related behaviours, which have a greater impact on the demand for health services and on the quality of life of people, or that address risk factors for NCDs are the most studied, owing to their influence on population morbidity and mortality. Thus, surveys have gained considerable importance in the assessment of lifestyle factors such as smoking, alcohol consumption, physical activity, sedentary behaviour, diet quality, overweight/obesity, hypertension, diabetes, violence, accidents, and mental disorders (Barros, 2008; Malta, Célia, & Szwarcwald, 2017).

The use of surveys including anthropometric indicators and food consumption are relevant in the assessment and monitoring of the dietary, nutritional, and health status conditions of the population (Sperandio & Priore, 2017), data widely described in the literature (Willet, 2013). In the case of dietary patterns, monitoring is an important mechanism to identify changes or trends in consumption and to understand the relationship between dietary exposure and varied health outcomes, as well as providing information relating to food, along with considering the nutritional interactions involved. The pattern of food consumption can also be used as an indirect indicator of nutritional status (Silva, Lyra, & Lima, 2016).

Some of the more common dietary surveys used in the identification of dietary patterns are the food record or diary, direct weighing of food, 24-hour dietary recall (R24h), dietary history, and the food frequency questionnaire (FFQ). There is no 'gold standard' method because each has its advantages and limitations. Therefore, the choice of method will depend on the characteristics of the population assessed and the objectives of the study (Fisberg, Marchioni, & Colucci, 2009; Shim, Oh, & Kim, 2014; Silva et al., 2016; Sperandio & Priore, 2017; Willet, 2013).

For anthropometric assessments, a direct method of assessment of nutritional status, population surveys have frequently utilized measurements of weight and height, owing to the ease of assessment and low cost of the equipment used, as well as the simplicity of the method, the use of non-invasive techniques, the reduced time to validate measures, the ease of training of personnel for data collection, and the reliability, when measured and evaluated properly by trained professionals (Duarte, 2007; Guedes, 2007; Sperandio & Priore, 2017).

Another advantage attributed to anthropometry is the easy comparison of measurements obtained with population and international standards used in the diagnosis of nutritional status, according to sex and age range, or the comparison between measurements, for the verification of changes in the dimensions and body composition of individuals (Nacif, 2007). On the other hand, the limitations of this method are related to their validation and accuracy, because it depends not only on the adequate training of field staff, but also on the use of calibrated equipment and standardization of the measures (Sperandio & Priore, 2017).

The reliability in the application of the method, the standardization of measures, and the possibility of international comparisons influence the reliability of the data and the reproducibility of the studies. Thus, this study aimed to perform a systematic review of population-based health and nutrition surveys, conducted with adults and the elderly in the Americas, Europe, and Oceania, to identify which anthropometric and food consumption methods were most frequently used, along with their applicability and limitations. These continents were selected because the dietary habits of Western populations are similar, in that there is a growing inclusion in the diet of foods high in calories, fat, and refined sugars, while they have low consumption of fruits and vegetables and fiber, in contrast to the diets in Eastern (Arab and Asian) and African countries.

2. Methodology

Design

This was a systematic review of original studies, prepared according to the recommendations of Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2009 (Moher et al., 2015), designed with the objective of guiding

the dissemination of systematic reviews and meta-analyses in health field. The review was registered in the international prospective register of systematic reviews (PROSPERO) under the number 2017: CRD42017071392.

Search strategy

A virtual search was performed in the LILACS, PubMed, and Scopus databases, for original articles published between 1997 and 2017. The following search strategy was used in PubMed: ('Health Surveys' OR 'Nutrition Surveys' OR 'Diet Surveys') AND ('Nutritional Status' OR 'Nutrition Assessment' OR 'Nutrition, Public Health') AND (adult OR adults OR aged OR elderly) AND ('cross-sectional studies' OR surveys) AND ('nationwide survey' OR 'national nutrition monitoring' OR 'national survey'). In the Scopus database, the key words used were: TITLE-ABS-KEY ('Health Surveys' OR 'Nutrition Surveys' OR 'Diet Surveys') AND TITLE-ABS-KEY ('Nutritional Status' OR 'Nutrition Assessment' OR 'Nutrition, Public Health') AND TITLE-ABS-KEY (adult OR adults OR aged OR elderly) AND TITLE-ABS-KEY ('cross-sectional studies' OR surveys) AND TITLE-ABS-KEY ('nationwide survey' OR 'national nutrition monitoring' OR 'national survey'). In LILACS the following combination of terms was used: (Inquérito OR pesquisa) AND (saúde OR nutrição OR dieta OR 'consumoalimentar' OR 'estadonutricional' OR antropometria OR "composição corporal") AND (nacional OR populacional). The searches were performed from April to July 2017 and confined to articles published between January 1997 and July 2017, in Portuguese, English, or Spanish. Data were analyzed in 2018.

Inclusion and exclusion criteria

The papers were considered eligible when they met the following inclusion criteria: 1) Observational studies; 2) Performed in adult and/or elderly populations; (3) Surveys carried out over the last 20 years (1997 to 2017); 4) Conducted in Europe, Americas, and/or Oceania; and 5) Available in Portuguese, English, or Spanish. Systematic reviews, meta-analyses and studies involving pregnant women were excluded from the study.

Synthesis and comparison of results

Initially, two evaluators independently reviewed the titles and abstracts to verify if those met the proposed eligibility criteria. Then, the complete texts of the papers were read. Subsequently, each evaluator completed, independently, a data mining spreadsheet, including, in addition to the evaluation of the methodological quality, the following items: 1) Characteristics of the studies: authors, year of publication, place of study (country and continent), and the evaluation of the methodological quality score. The population of each study was described according to the number and age of participants. 2) Characteristics of surveys: name, year, variables, method, frequency of application of the nutritional survey, and evaluation criteria of the anthropometric nutritional status and the respective cut-off points. Discrepancies in the evaluation were resolved through discussion between the evaluators, and, in case of doubt, a third reviewer was consulted.

Evaluation of the methodological quality of the studies

The methodological quality of the selected papers was evaluated and scored according to the recommendations of the Newcastle-Ottawa quality assessment scale (NOS) for cohort studies, and the Agency for Healthcare Research and Quality (AHRQ) checklist for cross-sectional studies. The NOS evaluation consists of eight questions, which include items such as: selection of participants, comparability between the subjects and verification of exposure. The papers are scored as 'Good', 'Adequate', or 'Poor', in accordance with the score received on each item of the scale (items are identified with one or no stars). The sum of these items (stars) classifies the article. The AHRQ consists of 11 items, with the options 'Yes', 'No', or 'Unclear'. A score '0' is attributed to items evaluated with 'No' or 'Unclear', and score '1' for those evaluated with 'Yes' (Shi,

Ying, Du, & Shen, 2017).

To better present the results, the score evaluated by NOS was converted into quality categories, based on the document from the AHRQ (AHRQ. Agency for Healthcare Research and Quality). Based on these results, three categories for evaluation were established: 0–3, 4–7, and 8–11, indicating low, moderate, and high quality, respectively.


Criteria for evaluation of methodological quality of primary studies

The following indicators of anthropometric evaluation and assessment of habitual consumption were considered as standard in the assessment of the methodological quality of the primary studies: 1) BMI: two or more measurements; 2) food surveys: at least two applications (for 24-hour dietary recall and food record), as recommended in the literature (Hoffmann et al., 2002). To check the quality of the anthropometric data collected, we investigated if the papers mentioned calibration of anthropometric instruments and training of the interviewers.

3. Results

The search recovered 615 documents. In addition, another 22 articles were added by manual search. 53 documents of the 637 were duplicated and for this reason were excluded, resulting in 584 abstracts. After reading the title and abstract, 253 articles were excluded. Thereafter, after considering the inclusion and exclusion criteria of 331 articles read in full, 48 studies were included in the review (Figure 1).

Figure 1. Flowchart of study selection.



F

Source: Authors.


Methodological quality and characteristics of the studies

Given the criteria for the classification of methodological quality, for the 48 studies included, 31 were classified as moderate quality (64.6%) and 17 as high quality (35.4%). Regarding the continent where the study was implemented, 45.8%


(n=22) were conducted in the Americas (Acosta, 2013; Acosta et al, 2005; Barquera et al., 2009, 2010; Barr, DiFrancesco, & Fulgoni, 2016; Bays, Chapman, & Grandy, 2007; Bezerra & Sichieri, 2011; Ferreira & Benicio, 2015; Gray-Donald, Jacobs-Starkey, & Johnson-Down, 2000; Guendelman et al., 2013; Kant, Graubard, 2015; Kordas et al., 2012; Melano-Carranza, Lasses Ojeda, & Ávila-Funes, 2008; Meller et al., 2014; Pavão, 2013; Pereira, Duffey, Sichieri, 2014; Pisabarro et al., 2009; Ponce et al., 2013; Sichieri et al., 2015; Tande, Magel, & Strand, 2009; Velasquez-Melendez et al., 2015), 47.9% (n=23) in Europe (Alkerwi et al., 2015; Alkerwi et al., 2015a; Beltrán-de-Miguel, Estévez-Santiago, & Olmedilla-Alonso, 2015; Bjerme et al., 2013; Cooper et al., 2014; Gazan et al., 2016; Gose et al., 2016; Kruizenza et al., 2003; Marcenes et al., 2003; Meier et al., 2010; Nissensohn et al., 2016, 2017; Pot et al., 2015; Pot et al, 2014; Prynne et al. , 2009; Ruiz et al., 2016; Tressou et al., 2016; Vernay et al., 2009; Whitton et al., 2011), and 6.3% (n=3) in Oceania(Bell, Edwards, & Grieger, 2015; Lei et al, 2016; Mohr et al., 2007). In relation to countries, Brazil contributed 14.6% (n=7) of the publications (Bezerra & Sichieri, 2011; Ferreira & Benicio, 2015; Meller et al., 2014; Pavão, 2013; Pereira, Duffey, Sichieri, 2014; Sichieri et al., 2015; Velasquez-Melendez et al., 2015); Spain (Beltrán-de-Miguel et al., 2015; Nissensohn et al., 2016, 2017; Ruiz et al., 2015, 2016), the United States (Bays et al., 2007; Cifelli et al, 2016; Guendelman et al., 2013; Kant, Graubard, 2015; Tande et al., 2009), and England (Cooper et al., 2014; Pot et al., 2015; Pot et al., 2014; Prynne et al., 2009; Whitton et al, 2011) contributed five each; Mexico (Barquera et al., 2009, 2010; Melano-Carranza et al., 2008; Ponce et al., 2013) four; Australia (Bell et al., 2015; Lei et al., 2016; Mohr et al., 2007) and France(Gazan et al., 2016; Tressou et al., 2016; Vernay et al., 2009) three; Canada (Barr et al., 2016; Gray-Donald et al., 2000), Colombia (Acosta, 2013; Kordas et al., 2012), Luxembourg (Alkerwi et al., 2015; Alkerwi et al., 2015a) and Netherlands (Kruizenza et al, 2003; Sluik et al, 2014) two; Germany (Gose et al., 2016), Cuba (Acosta et al, 2005), the United Kingdom (Marcenes et al., 2003), Sweden (Bjerme et al, 2013), Switzerland (Meier et al., 2010), and Uruguay (Pisabarro et al., 2009) contributed one study each (Figure 2).

The studied population ranged from 273 (Bjerme et al, 2013) to 215,354 (Bays et al., 2007) individuals. In relation to the nutrition variables analysed, 35.4% (n=17) of the papers evaluated food consumption and anthropometry, 35.4% (n=17) only food consumption, and 29.2% (n=14) only anthropometry (Figure 2).

Figure 2. Characteristics of studies.



a. Contribution of countries in health and nutrition studies.



b. Methods used for assessment of food consumption and anthropometry.

Source: Authors.

The main population surveys on nutrition and health that served as the basis for the papers, alone or in comparison with other surveys, were: the National Health and Nutrition Examination Survey (NHANES; USA) (Bays et al., 2007; Cifelli et al., 2016; Guendelman et al., 2013; Kant; Graubard, 2015; Tande et al., 2009), with five derived studies; Anthropometry, Intake and Energy Balance in Spain (ANIBES; Spain) (Nissensohn et al., 2016, 2017; Ruiz et al., 2015, 2016) and National Survey of Health and Development (NSH; England) (Cooper et al., 2014; Pot et al., 2014; Pot et al., 2015; Prynne et al., 2009) with four articles each; Italian National Food Consumption Survey (INRAN-SCAI; Italy) (Leclercq et al., 2009; Nissensohn et al., 2017; Sette et al., 2018), Brazilian Household Budget Survey/Pesquisa de Orçamentos Familiares (Brazil) (Bezerra & Sichieri, 2011; Pereira, Duffey, Sichieri, 2014; Sichieri et al., 2015) with three publications; National Demography and Health Survey/Pesquisa Nacional de Demografia e Saúde (PNDS; Brazil) (Meller et al., 2014; Pavão, 2013), Encuesta Nacional de Salud y Nutrición (ENSANUT; Mexico) (Barquera et al., 2010; Ponce et al., 2013), Encuesta Nacional de Situación Nutricional (ENSIN; Colombia) (Acosta, 2013; Kordas et al., 2012), Individual and national study on food consumption (INCA 2; France) (Gazan et al., 2016; Tressou et al., 2016), British National Diet and Nutrition Survey (NDNS; UK/England) (Marcenes et al., 2003; Whitton et al., 2011) and Observation of Cardiovascular Risk Factors in Luxembourg

(ORISCAV-LUX; Luxembourg)(Alkerwi et al, 2015; Alkerwi et al., 2015a) with two publications each. The following served as a basis for one publication each: Dutch National Food Consumption Survey (DNFCS; Netherlands) (Sluik et al, 2014), NutriNet-Santé Dataset (France) (Nissensohn et al., 2017), ENDS (Colombia) (Acosta, 2013), Australian Health Survey (Australia) (Bell et al., 2015), Estudio Nacional de Salud y Envejecimiento en México (ENASEM; Mexico) (Melano-Carranza et al., 2008), Encuesta Nacional de Ingesta Dietética (ENIDE; Spain)(Beltrán-de-Miguel et al., 2015), Étude Nationale Le Nutrition Santé(ENNS; France) (Vernay et al., 2009), Segunda Encuesta Nacional de Sobrepeso y Obesidad (ENSO 2; Uruguay) (Pisabarro et al., 2009), Food Habits of Canadians Survey (Canada) (Gray-Donald et al., 2000), Mexican National Health and Nutrition Survey 2006 (Mexico) (Barquera et al., 2009), National Screening on Malnutrition (Netherlands) (Kruizenga HM, Wierdsma NJ, Vanbokhorst MAE, De Vander Schueren, Hollander HJ, 2003), Nielsen Media Research (Australia) (Mohr et al., 2007), National Nutrition and Physical Activity Survey (NNPAS; Australia) (Lei et al., 2016), German National Nutrition Survey (NVS II; Germany) (Gose et al., 2016), German National Nutrition Monitoring (NEMONIT;Germany) (Gose et al., 2016), National Health Research/Pesquisa Nacional de Saúde(PNS;Brazil) (Velasquez-Melendez et al., 2015), Research Social Dimension of Inequalities/Pesquisa Dimensão Social das Desigualdades(PDSD; Brazil) (Pavão, 2013), Segunda Encuesta Nacional sobre Factores de Riesgo y Afecciones Crónicas no Transmisibles de la Población Cubana (Cuba) (Acosta et al, 2006), Study to Help Improve Early Evaluation and Management of Risk Factors Leading to Diabetes (SHIELD;USA)(Bays et al., 2007), The National Survey Riksmaten (Sweden) (Bjermo et al, 2013) and The Swiss Health Survey (Switzerland)(Meier et al., 2010).

Methods used for assessment of food consumption and anthropometry

Food record was the more frequent survey method, used in 16 studies (Bezerra & Sichieri, 2011; Bjermo et al, 2013; Gazan et al., 2016; Leclercq et al., 2009; Marcenes et al., 2003; Nissensohn et al., 2017; Pereira, Duffey, Sichieri, 2014; Pot et al., 2014; Pot et al., 2015; Prynne et al., 2009; Ruiz et al., 2016, 2015; Sette et al., 2018; Sichieri et al., 2015; Tressou et al., 2016; Whitton et al, 2011), followed by the R24h, used in 11 studies (Barr et al., 2016; Bell et al., 2015; Beltrán-de-Miguel et al., 2015; Cifelli et al., 2016; Gose et al., 2016; Kant, Graubard, 2015; Lei et al., 2016; Pisabarro et al., 2009; Sluik et al, 2014; Tande et al., 2009; Vernay et al., 2009), the FFQ, included in seven articles (Alkerwi et al., 2015; Alkerwi et al., 2015a; Bjermo et al, 2013; Meier et al., 2010; Mohr et al., 2007; Ponce et al., 2013; Velasquez-Melendez et al., 2015) and, finally, the food diary, applied in three studies (Nissensohn et al., 2016, 2017; Vernay et al., 2009). The application frequency varied according to the type of food survey conducted. The number of food records ranged from two to seven days, the R24h was applied from one to five times, and the food diary was used for three days, in all studies.

Nine of the 14 articles that used the food record performed four or more assessments (Bezerra & Sichieri, 2011; Bjermo et al, 2013; Gazan et al., 2016; Marcenes et al., 2003; Pot et al., 2014; Pot et al., 2015; Prynne et al., 2009; Tressou et al., 2016; Whitton et al., 2011). Along with the R24h, 11 papers applied this method, at least two applications were performed in at least seven studies (Bell et al., 2015; Cifelli et al., 2016; Gose et al., 2016; Gray-Donald et al., 2000; Lei et al., 2016; Melano-Carranza et al., 2008; Sluik et al., 2014), as recommended by the National Cancer Institute (2015). The evaluation of habitual consumption, dietary pattern, ingestion of foods or food groups, energy, and macro and/or micronutrients, were the purpose of the application of the R24h in nine studies (Bell et al., 2015; Beltrán-de-Miguel et al., 2015; Cifelli et al., 2016; Gose et al., 2016; Gray-Donald et al., 2000; Kant, Graubard, 2015; Lei et al., 2016; Sluik et al., 2014; Tande et al., 2009). The R24h was used to evaluate consumption of alcohol and breakfast (being a consumer or not) in two other papers (Barr et al., 2016; Vernay et al., 2009).

In relation to anthropometry, BMI was the indicator used in all 30 studies designed to assess anthropometric nutritional status(Acosta, 2013; Acosta et al, 2006; Barquera et al., 2009, 2010; Barr et al., 2016; Bays et al., 2007; Bell et al.,

2015; Bezerra & Sichieri, 2011; Cooper et al., 2014; Ferreira & Benicio, 2015; Guendelman et al., 2013; Kant, Graubard, 2015; Kordas et al., 2012; Kruizenga et al, 2003; Leclercq et al., 2009; Lei et al., 2016; Marcenes et al., 2003; Meier et al., 2010; Melano-Carranza et al., 2008; Meller et al., 2014; Nissensohn et al., 2017; Pavão, 2013; Pisabarro et al., 2009; Ponce et al., 2013; Pot et al., 2015; Sette et al., 2018; Sichieri et al., 2015; Sluik et al., 2014; Velasquez-Melendez et al., 2015; Vernay et al., 2009), with the exception of the study of Tande et al.(Tande et al., 2009), which evaluated only waist circumference. Some of the studies also evaluated other body measurements, such as waist circumference (Acosta et al, 2006; Barquera et al., 2009; Bell et al., 2015; Meller et al., 2014; Nissensohn et al., 2017) and waist-hip ratio (Ferreira & Benício, 2015). About the BMI evaluation, 20 papers used the classification proposed by the World Health Organization (WHO) (Acosta, 2013; Acosta et al, 2006; Barquera et al., 2009, 2010; Barr et al., 2016; Bays et al., 2007; Bell et al., 2015; Bezerra & Sichieri, 2011; Cooper et al., 2014; Ferreira & Benicio, 2015; Guendelman et al., 2013; Kant ; Graubard, 2015; Kordas et al., 2012; Kruizenga et al, 2003; Lei et al., 2016; Marcenes et al., 2003; Meier et al., 2010; Melano-Carranza et al., 2008; Meller et al., 2014; Nissensohn et al., 2017; Pavão, 2013; Pisabarro et al., 2009; Ponce et al., 2013; Pot et al., 2015; Sichieri et al., 2015; Velasquez-Melendez et al., 2015; Vernay et al., 2009). The studies of Melano-Carranza et al. (Melano-Carranza et al., 2008), Nissensohn et al. (Nissensohn et al., 2017), Pot et al. (Pot et al., 2015), Cooper et al. (Cooper et al., 2014), Sluik et al. (Sluik et al, 2014), Sette et al. (Sette et al., 2018), and Leclercq et al. (Leclercq et al., 2009) only reported the mean BMI values in the population studied. Acosta et al. (Acosta SJ, Sánchez et al, 2006) used the classification of the Food and Agriculture Organization of the United Nations (Shetty & James, 1994). The study by Barr et al. (Barr et al., 2016) did not describe the classification used. Other studies used classifications of other international institutions such as the International Obesity Task Force (Pisabarro et al., 2009) and the National Institutes of Health Clinical Guidelines (Bays et al., 2007).

With regard to the quality of the measures for composition of BMI, 11 papers reported that interviewers/researchers were trained to perform the measurements (Barquera et al., 2009, 2010; Barr et al., 2016; Cooper et al., 2014; Guendelman et al., 2013; Kordas et al., 2012; Meller et al., 2014; Pisabarro et al., 2009; Ponce et al., 2013; Pot et al., 2015; Vernay et al., 2009), four studies reported the calibration of equipment (Acosta SJ, Sánchez et al, 2006; Kordas et al., 2012; Meller et al., 2014; Ponce et al., 2013), and only two articles (Ferreira & Benicio, 2015; Meller et al., 2014) reported performing two repetitions of each measurement of weight and height, as standardized by the WHO for collection of these data. Only the study of Meller et al. (Meller et al., 2014) reported all the information about the survey (training of the interviewers, calibration of equipment, and replication of anthropometric measures).

Table 1. Characteristics of the studies and population-based surveys on health and nutrition, performed in the Americas, Europe, and Oceania, from 1997 to 2017.

Characteristics of the Studies								Characteristics of surveys			
Author/Year	Continent	Country	Study Design	n	Main	Age range (years)	Score/ Methodological quality	Survey	Variables	BMI cut-off points	Nº Records
Acosta K., 2013.	Americas	Colombia	Cross-sectional	150,733	To calculate obesity concentration indices among adults in Colombia.	18-64	7 (moderate)	ENDS (2005) and ENSIN (2010)	BMI	WHO, 1995	-
Acosta S. et al., 2005.	Americas	Cuba	Cross-sectional	19,519	To evaluate the nutritional status of the resident population in the urban area of Cuba and compare evolutionarily the changes experienced in relation to the previous survey.	>20	6 (moderate)	Segunda Encuesta Nacional sobre Factores de Riesgo y Afecciones Crónicas no Transmisibles de la Población Cubana (2000-2001)	BMI CC CQ	FAO, 1994 WHO, 1998	-
Alkerwi A. et al, 2015.	Europe	Luxembourg	Cross-sectional	1,352	To examine the association between nutritional awareness and diet quality, as indicated by energy density, dietary diversity and adequacy to achieve dietary recommendations, while considering the potentially important role of socioeconomic status	18-69	7 (moderate)	ORISCAV-LUX (2007-2008)	Food consumption	-	Semi-quantitative FFQ
Alkerwi A.	Europe	Luxembourg	Cross-	1,351	To compare the ability of five diet quality indices,	18-69	7 (moderate)	ORISCAV-LUX (2007-	Food	-	QFC

et al, 2015a.			g	sectional		namely the Recommendation Compliance Index, Diet Quality Index-International, Dietary Approaches to Stop Hypertension, Mediterranean Diet Score, and Dietary Inflammatory Index, to detect changes in chronic disease risk biomarkers.			2009)	consumption		
Barquera S. et al, 2009.	Americas	Mexico		Cross-sectional	33,023	To estimate the prevalence of overweight, obesity and central adiposity in Mexico, and to explore trends compared to the previous Mexican National Health Survey and to Mexican-Americans.	>20	8 (high)	Mexican National Health and Nutrition Survey (2006)	BMI CC	WHO, 1998	-
Barquera S. et al, 2010.	Americas	Mexico		Cross-sectional	14,630	To describe the prevalence of hypertension among Mexican adults, and to compare to that observed among Mexican-Americans living in the US.	>20	8 (high)	ENSANUT (2006)	BMI	WHO, 1998	-
Barr S. et al, 2016.	Americas	Canada		Cross-sectional	12,337	To examine the association of breakfast consumption, and the type of breakfast consumed, with body mass index and prevalence rates and odds ratios of overweight/obesity among Canadian adults.	>18	7 (moderate)	Canadian Community Health Survey Cycle 2.2 (2004)	Dietary intake BMI	BMI > 25.0 for definition of overweight/obesity in all age groups	R24h 1 day

Bays H. et al, 2007.	Americas	USA	Cross-sectional	215,354	To explore the relation between body mass index and prevalence of diabetes mellitus, hypertension and dyslipidaemia; examine BMI distributions among patients with these conditions; and compare results from two national surveys.	>18	7 (moderate)	SHIELD (2004) NHANES (1999-2002)	BMI	Adaptation of the 1998 National Heart, Lung and Blood Institute - National Institutes of Health Clinical Guidelines.	-
Bell L. et al, 2015.	Oceania	Australia	Cross-sectional	2,415	To identify dietary patterns in Australian adults, and to determine whether these dietary patterns are associated with metabolic phenotype and obesity.	>45	6 (moderate)	Australian Health Survey (2011-2013)	BMI CC Food consumption	WHO, 1998	R24h 2 days
Beltrán-de-Miguel B. et al, 2015.	Europe	Spain	Cross-sectional	3,000	To assess the intake of the individual components of vitamin A and major dietary sources in the Spaniards using data on food consumption from Spanish National Dietary Intake Survey (2009–2010).	18-64	6 (moderate)	ENIDE (2009-2010)	Consumption of vitamin A		R24h 1 day 3-day Food Diary
Bezerra I; Schieri R., 2011.	Americas	Brazil	Cross-sectional	33,393	To evaluate whether a diversity of healthy foods in a household would decrease the availability of unhealthy foods and to evaluate the association between a healthy dietary diversity score and nutritional	20-65	8 (high)	Brazilian Household Budget Survey (2002-2003)	BMI Food consumption	WHO, 1998	7-day Food Record

status among adults.												
Bjermo H. et al, 2013.	Europe	Sweden	Cross-sectional	273	To examine the body burden of lead, mercury, and cadmium in blood among Swedish adults and the association between blood levels, diet and other lifestyle factors.	18-80	8 (high)	The National Survey Riksmaten (2010-2011)	Food and dietary consumption	-	4-day food record	FFQ foods not widely consumed
Cifelli C. et al, 2016.	Americas	USA	Cross-sectional	17,387*	To use national survey data to model different dietary scenarios to assess the potential effects of increasing plant-based foods (and concomitantly decreasing animal foods) or dairy foods on macronutrient intake and nutrient adequacy.	2-18 and >19	7 (moderate)	NHANES (2007-2010)	Food and dietary consumption	-	R24h 2 days (in person + telephone contact)	
Cooper R. et al, 2014.	Europe	England	Cross-sectional	2,229* (1,511 adults/elderly)	To examine the associations of body mass index from age 15 years onwards with low muscle mass, strength, and quality in early old age.	15, 20, 26, 36, 43, 53, 60-64	6 (moderate)	NSHD (data from 1946 cohort)	BMI	Mean BMI	-	
Ferreira R.; Benício M, 2015.	Americas	Brazil	Cross-sectional	11,961	To determine the influence of reproductive history on the prevalence of obesity in Brazilian women and effect of socioeconomic variables on the association between parity and excess weight.	20-49	6 (moderate)	PNDS (2006)	BMI	WHO, 2000	-	
Gazan R. et	Europe	France	Cross-	1,918	To examine the association between	>18	7 (moderate)	INCA 2	Consumption of food,	-	7-day Food	

al, 2016.			sectional		drinking water intake and diet quality, and to analyse the adherence of French men and women to the European Food Safety Authority 2010 Adequate Intake.				(2005-2007)	drinks, and water		Record
Gose M. et al, 2016.	Europe	Germany	Cohort	1,840*	To assess changes in food consumption and nutrient intake in Germany.	14-65 and >65		8 (high)	NVS II (2005/2007)	Dietary consumption	-	R24h 2 days
Gray-Donald K. et al, 2000.	Americas	Canada	Cross-sectional	1,722* (1,544 adults/elderly)	To monitor whether changes in dietary intake have occurred since the last Canadian dietary survey, conducted a generation ago (1970).	13-17 and 18-65		7 (moderate)	Food Habit of Canadians survey (1997-1998)	Food consumption	-	R24h - 1 day (repeated in 30% of the sample) with the use of models of food portions.
Guendelman S. et al, 2013.	Americas	USA	Cross-sectional	979	To examine actual and perceived weight in national cohorts of Mexican-origin adult men in Mexico and the United States (US).	20-59		7 (moderate)	NHANES (2001/2006) and ENSANUT (2006)	Calculated BMI and self-perceived BMI	WHO, 1998	-
Kant A. et al, 2015.	Americas	USA	Cross-sectional	62,298	To examine time trends in the distribution of day's intake into individual meal and snack behaviors and related attributes in the United States adult population.	20-74		9 (high)	NHANES (1971-74, 1976-1980, 1988-1994, 1999-2000, 2001-2002, 2003-2004, 2005-2006, 2007-2008,	Food consumption of meals and snacks BMI	WHO, 1998	R24h 1 day (in paper form, interview by telephone, computer)

									and 2009–2010)			
Kordas K. et al, 2013.	Americas	Colombia	Cross-sectional	3,267* (2,612 adults)	To investigate the association among iron deficiency (ID), anemia, and weight status among nonpregnant Colombian females aged 13–49 y.	13-17 and 18-49	7 (moderate)		ENSIN (2005)	BMI	WHO, 1998	
Kruizenga H. et al, 2003.	Europe	Netherlands	Cross-sectional	7,606	To determine the prevalence of disease-related malnutrition in The Netherlands in all fields of medical care and to investigate the involvement of the dietitian in the treatment of malnutrition.	>18	8 (high)		National Screening on Malnutrition (2001)	BMI	WHO, 1998	-
Leclercq C. et al, 2009.	Europe	Italy	Cross-sectional	2,831	To present the main results of the Italian National Food Consumption Survey INRAN-SCAI 2005–06.	>18	8 (high)		INRAN-SCAI (2005-2006)	Food consumption BMI	Mean BMI	self-administered 3-day food record
Lei L. et al, 2016.	Oceania	Australia	Cross-sectional	8,202* (6,326 adults and elderly)	To examine the AS and free sugar intakes and the main food sources of AS among Australians.	>02	8 (high)		NNPAS (2011-2012)	BMI Food consumption	WHO, 1998	R24h 2 days (in person + telephone contact)
Marcenes W. et al, 2003.	Europe	UK	Cross-sectional	949	To review the major findings from a large representative and comprehensive national survey in Great Britain to which the numbers of teeth and dentures affected older people's ease of eating, nutrient	>65	6 (moderate)		NDNS (1998)	BMI Food and dietary consumption	WHO, 1998	4-day Heavy food record

					intake, nutritional status, and body mass index.								
Meier M. et al, 2010.	Europe	Switzerland	Cross-sectional	1,786	To assess whether Swiss residents aged 15–24 years follow current nutritional guidelines and whether differences exist according to gender and weight status.	15-24	9 (high)	The Swiss Health Survey (2007)	BMI Food consumption	WHO, 1998	FFQ		
Melano-Carranza E., 2008	Americas	Mexico	Cross-sectional	2,029	To determine factors associated with failure to adhere to treatment for diagnosed hypertension among a representative sample of older Mexican adults living in the community.	>65	7 (moderate)	ENASEM (2001)	BMI	Mean BMI	-		
Meller F. et al, 2014.	Americas	Brazil	Cross-sectional	14,101	To evaluate the association between waist circumference and body mass index of Brazilian women of childbearing age studied in the National Demographic and Health Survey, in 2006.	18-49	6 (moderate)	PNDS (2006)	BMI/CC	WHO, 1998	-		
Mohr P. et al, 2007.	Oceania	Australia	Cross-sectional	20,527*	To identify key predictors of fast-food consumption from a range of demographic, attitudinal, personality and lifestyle variables.	>14	6 (moderate)	Nielsen Media Research (2004-2005)	Food consumption	-	FFQ fast food		

Nissensohn M. et al, 2016.	Europe	Spain	Cross-sectional	2,007* (1,784 adults/elderly)	To quantify the total water and beverage intake, and to explore associations between the types of beverage consumed and energy intake.	9-75	8 (high)	ANIBES (2013)	Consumption of foods and beverages	-	3-days Food Record (food diary) (using a digital camera, tablet, interview by phone, or form)
Nissensohn M. et al, 2017.	Europe	Spain	Cross-sectional	99,111	To compare the average daily consumption of foods and beverages in adults of selective samples of the European Union population to understand the contribution of these to the total water intake, evaluate if the EU adult population consumes adequate amounts of total water according to the current guidelines, and to illustrate the real water intake in Europe.	18-75	6 (moderate)	ANIBES (Spain 2013); INRAN-SCAI Dataset (Italy – 2005-2006); NutriNet-Santé Dataset (France – 2009-2010)	BMI - CC Consumption of foods and beverages	Mean BMI	Italy - semi-structured 3-day diary France - 3-day food record Spain - 3-day food record
Pavão A. et al, 2013.	Americas	Brazil	Cross-sectional	12,324	To investigate the association between self-rated health and social and demographic factors, health behavior, and morbidity.	>20	7 (moderate)	PDS (2008)	BMI	WHO, 1998	-
Pereira R. et al, 2014.	Americas	Brazil	Cross-sectional	34,003* (26,522)	To examine the patterns of consumption of foods high in solid fats and added sugars in the first	>10	6 (moderate)	Brazilian Household Budget Survey	Consumption of saturated and trans fats	-	2-day Food Record

				adults/ The elderly)	Brazilian nationally representative individual dietary survey.			(2008-2009)	Consumption of added sugar			
Pisabarro R. et al, 2009.	Americas	Uruguay	Cross-sectional	900	To evaluate prevalence of obesity, its comorbidities and predisposing factors through the Second National Survey on Overweight and Obesity in Uruguay in people between 18 and 65 years, or older, carried out in 2006.	18-65 and >65	6 (moderate)	ENSO 2 (2006)	BMI	International Obesity Force	-	Task
Ponce X. et al, 2013.	Americas	Mexico	Cross-sectional	15,675	To characterize the health and nutritional status of the Mexican population in all age groups.	19-59	8 (high)	ENSANUT (2006)	BMI Food consumption	WHO, 1998		Semi-quantitative 7-day FFQ
Pot G. et al, 2014.	Europe	England	Cohort	989	To describe changes in food consumption patterns and food availability in an ageing population.	36, 43, 53, 60-64	8 (high)	NSHD (cohort 1946)	Food consumption	-		Pre-structured 5-day food record
Pot G. et al, 2015.	Europe	England	Cross-sectional	1,768	To study associations between irregular consumption of energy intake in meals and cardio-metabolic risk factors.	53	8 (high)	NSHD (cohort of 1946 - the year 1999 was used for this study).	BMI Dietary Intake	Mean BMI		5-day food record
Prynn C. et al, 2009.	Europe	England	Cohort	4,028	To quantify more precisely the meat intake of a cohort of adults in the	43 and 53	8 (high)	NSHD (cohort of	Ingestion of meat	-		Pre-structured 5-day food

					UK by disaggregating composite meat dishes.			1946)		record	
Ruiz E. et al, 2015	Europe	Spain	Cross-sectional	2,009* (1,861 adults/elderly)	To contribute to updating data of dietary energy intake and its main sources from food and beverages, according to gender and age.	9-75	6 (moderate)	ANIBES (2013)	Consumption of energy and nutrients Food consumption	-	3-day food record (A three-day dietary record) (with the use of digital camera, tablet, interview by phone, or form)
Ruiz E. et al, 2016.	Europe	Spain	Cross-sectional	2,009* (1,861 adults/elderly)	To analyze dietary macronutrient intake and its main sources according to sex and age.	9-75	6 (moderate)	ANIBES (2013)	Consumption of foods and beverages Dietary Intake	-	3-day Food Record
Sette S. et al, 2011.	Europe	Italy	Cross-sectional	2,830	To describe energy and nutrient intakes in Italy.	>18	7 (moderate)	INRAN-SCAI (2005-2006)	Energy, macro, and micronutrients Alcohol intake BMI	Mean BMI	self-administered 3-day food record
Sichieri R. et al, 2015.	Americas	Brazil	Cross-sectional	34,003* (26,862 adults/elderly)	To identify major food group contributors of energy intake.	>10	9 (high)	Brazilian Household Budget Survey (2008-2009)	BMI >25kg/m ² Food consumption	WHO, 1998	2-day Food Record
Sluik D. et al, 2014.	Europe	Netherlands	Cross-sectional	2100	To investigate associations between alcoholic beverage preference and dietary	19-69	7 (moderate)	DNFCS	Energy intake Frequency and absolute consumption	Mean BMI	R24h 2 non-consecutive days

					intake in The Netherlands.						of alcohol		
											BMI		
Tande D. et al, 2009.	Americas	USA	Cross-sectional	15,658	To describe relationships between the Health Eating Index and abdominal obesity among adults.	>20	7 (moderate)			NHANES III (1988-1994)	CC Index of healthy eating	-	R24h 1 day
Tressou J. et al, 2016.	Europe	France	Cross-sectional	2,624	To explore in details the fatty acids intakes in French adults using the most recent available data.	Adults	7 (moderate)			INCA 2 (2006-2007)	Consumption of fish or margarine	-	7-day Food Record
Velasquez-Melendez G.; Mendes M, 2015.	Americas	Brazil	Cross-sectional	34,362	To evaluate the prevalence of cardiovascular health in the Brazilian population.	>18	9 (high)			PNS (2013)	BMI Food consumption	WHO, 1998	Markers of food consumption questionnaire
Vernay M. et al, 2009.	Europe	France	Cross-sectional	3,115	To describe disparities in the prevalence of overweight and obesity across socioeconomic status groups in 18-74 year-old French adults.	18-74	7 (moderate)			ENNS (2006-2007)	Food consumption BMI Cardiovascular risk	WHO, 1998	R24h 3 days
Whitton C. et al, 2011.	Europe	England	Cross-sectional	4,321* (2,158 adults and elderly)	To report dietary intakes and main food sources of fat and fatty acids from the first year of the National Diet and Nutrition Survey rolling programme in the UK.	4-64	7 (moderate)			NDNS 2008-2009	Food and dietary consumption	-	4-day food record

*studies that included children and/or adolescents, besides adults and the elderly. Source: Authors.

4. Discussion

Food consumption surveys are used to collect information about the preparation and consumption of food, through observations by skilled personnel (Shim et al., 2014). In the present analysis, the food record was the most frequently used nutritional survey to assess food consumption in large population surveys, followed by the R24H. Some countries have used this type of dietary inquiry, including Brazil (Bezerra & Sichieri, 2011; Pereira; Duffey; Sichieri, 2014; Sichieri et al., 2015), as well as developed countries such as Spain (Nissensohn et al., 2017; Ruiz et al., 2015, 2016), England (Pot et al., 2015; Pot et al., 2014; Prynne et al., 2009; Whitton et al., 2011), France (Gazan et al., 2016; Tressou et al., 2016), the United Kingdom (Marcenes et al., 2003), and Switzerland (Bjerme et al., 2013).

Although allowing a detailed assessment of food consumption, the food record dispenses with the interviewer and does not feature respondent bias. However, it does present some limitations such as the requirement for literacy and high motivation (Shim et al., 2014). It is possible that preference for its use in European countries is due to the easier application since an interviewer is not necessary. In this sense, it reduces the costs of the research. It should be emphasized that use of this method has greater reproducibility in populations with a high level of education.

Wide variations in the type of equipment used for data collection were observed in these studies, such as the use of tablets, digital cameras, photographic recordings, telephone interviews, and printed forms, which require a certain degree of skill for their use. It is also interesting that most papers that used the food record applied it for four or more days (Bezerra & Sichieri, 2011; Bjerme et al., 2013; Gazan et al., 2016; Marcenes et al., 2003; Pot et al., 2014; Pot et al., 2015; Prynne et al., 2009; Tressou et al., 2016; Whitton et al., 2011). Accordingly, the choice of this method over a long period of time requires some skills from the interwire, such as: good collaborative ability, motivation, and an understanding of the importance of the study. These avoid possible biases resulting from delay in the implementation of the method, such as underreporting or overestimation, as well as withdrawal of the interviewer (FAO, 2018).

The R24h also provides details of food intake and offers greater convenience to the respondent than food record. However, there is a possibility of both respondent and interviewer bias, and the method requires trained interviewers. In this review, only four articles described the training of the interviewers (Barr et al., 2016; Bell et al., 2015; Beltrán-de-Miguel et al., 2015; Kant; Graubard, 2015). As all the papers were based on national surveys, generally using secondary data from databases, it is assumed there was a lack of information on interviewer training. Furthermore, the information was not available in the database, but was described in the survey methodology.

Both the food record and the R24h require multiple days of evaluation to estimate habitual intake (Shim et al., 2014). Recording of several days of consumption when using short-term records, such as the R24h and the food record, is necessary to remove intrapersonal variability, reducing the random error inherent in usual food consumption (Morimoto et al., 2011).

In relation to anthropometric measurements, most of the studies used the BMI as an indicator to assess anthropometric nutritional status. The evaluation of body composition may be relevant as it evaluates the role of body components to health, as well as its relations with the emergence of NCDs. There are various methods for the assessment of body composition that are considered accurate and sophisticated, such as hydrostatic weighing and dual-energy x-ray absorptiometry (DEXA). However, their use in epidemiological studies is impractical because of the high cost. Therefore, BMI and waist circumference are widely used for anthropometric indicators in population studies due to their practicality and low cost (Rezende et al., 2007) .

In practice, the anthropometric data are compared for the interpretation of anthropometric nutritional status to reference values, which were obtained from specific populations (De Onis, 1996). Thus, it is necessary to consider different cut-off points for the interpretation of distinct life cycles, such as aging.

The WHO (WHO, 2002) defines an aged or 'older person' as a 60 years old individual or older. This definition, according to the WHO, is appropriate for developing countries; however, in some contexts, especially in developed countries, considering the significant increase in life expectancy in recent years, the classification of 65 years or more for an elderly person may be more appropriate.

However, it's considering that chronological age is not necessarily a precise marker in the monitoring of changes related to aging, because there are important differences related to health, participation, and levels of interdependence among people of the same age. Therefore, it is worth emphasizing that for the purpose of formulating public policies (one of the purposes of national surveys), these variations among older people should be considered (World Health Organization, 2005).

Despite its wide dissemination in epidemiological studies, the use of BMI for assessment of anthropometric nutritional status of the elderly has been questioned, owing to the changes in body composition resulting from the aging process (Cook et al, 2005). Thus, the employment of the same criteria for the classification of BMI in the general adult population and elderly patients is controversial, considering the reduced stature, accumulation of adipose tissue, and reduction of lean body mass and water in the body that occurs in the elderly individual (Silveira, Kac, & Barbosa, 2009).

The criteria of the WHO (<18,5 kg/m² – Underweight; 18,5 to 24,9 kg/m² - Normal weight ; 25,0 to 29,9 kg/m² - Pre-obesity; >30,0 kg/m²- obesity)(World Health Organization, 2000) and those proposed by Lipschitz (< 22 kg/m² - Underweight; 22,0 to 27,0 - Normal weight; < 27 kg/m² - Overweight) (Lipschitz, 1994) et al. are the most commonly used for the assessment of anthropometric nutritional status of the elderly.

In this systematic review, which aimed to evaluate the population studies conducted with adults and elderly patients, it was observed that in the majority of studies, the cut-off points of BMI classification proposed by the WHO for adults were also used for the elderly (Acosta, 2013; Barquera et al., 2009, 2010; Bell et al., 2015; Bezerra & Sichieri, 2011; Kantet al, 2003; Lei et al., 2016; Marcenes et al., 2003; Pavão, 2013; Sichieri et al., 2015; Velasquez-Melendez et al., 2015; Vernay et al., 2009). However, unlike the cut-off points proposed by Lipschitz et al (Lipschitz, 1994), the WHO classification does not consider the changes in body composition of the elderly. Therefore, it is necessary to critically analyse the classifications of BMI in large population surveys and consider the use of other anthropometric indicators for a complementary diagnosis or update of the criteria proposed by the WHO.

Thirty-five of the 45 papers performed anthropometric measurements. Information on the techniques and protocols used, interviewer training, and replication of anthropometric measures were available in most of these studies or in previous publications. However, in 40.0% (n = 14) (Acosta, 2013; Acosta et al, 2006; Barquera et al., 2009; Barros, 2008; Bell et al., 2015; Bjeremo et al, 2013; Lei et al., 2016; Marcenes et al., 2003; Meier et al., 2010; Pavão, 2013; Pisabarro et al., 2009; Pot et al., 2015; Sichieri et al., 2015; Tressou et al., 2016; Vernay et al., 2009), a lack of clarity was observed in the equipment calibration information, which did not appear in previous publications, nor even in official survey sites or documents investigated.

This review contains some limitations that must be considered, such as the possible non-inclusion of all health and nutrition surveys conducted on the continents surveyed, because not all databases were consulted. However, to circumvent this limitation, a manual search was performed to obtain a greater number of studies for inclusion.

It is worth mentioning this study included articles of high methodological quality; thus, the results presented here depict the more common methodologies used and accepted in the scientific community for assessment of food intake and anthropometric nutritional status, allowing, however, their critical analysis, to improve the reproducibility of health and nutrition studies.

5. Conclusion

This review identified the food record and BMI as the most common indicators of the evaluation of food consumption and anthropometry in population surveys conducted with adults and the elderly. The food record and the 24-hour dietary recall were the preferred methods for studies in developed countries, where high education levels and motivation of residents facilitate the application of these methods. BMI, owing to the ease of obtaining its component measurements and the wide application of its use, was the method of choice of the studies that performed anthropometric evaluation. Nevertheless, most of the studies did not use different classifications in the BMI assessment of adults and the elderly, disregarding the specificities of the changes in body composition and the physiological process of aging widely reported in the literature. The lack of clarity or omission in relation to information regarding the quality of dietary or anthropometric surveys, such as interviewer training, calibration of equipment, or replication of anthropometric measurements, were also issues observed in this review that should be reported.

Hence, it should be emphasized that, in population studies that assess health and nutrition, it is important to devise a proper study design, by selecting the method of food survey that best fits the objectives, the use of anthropometric indicators feasible in epidemiology, and the use of cut-off points appropriate for the population studied, with a view toward reducing the biases and providing valid data. Population surveys can thus provide reliable guidance for the formulation of public policies consistent with the epidemiological profile identified in these studies.

Abbreviations

PROSPERO: International prospective register of systematic reviews

PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses

NOS: Newcastle-Ottawa quality assessment scale

AHRQ: Agency for Healthcare Research and Quality

BMI: Body Mass Index

NHANES: National Health and Nutrition Examination Survey

ANIBES: Anthropometry, Intake and Energy Balance in Spain

NSH: National Survey of Health and Development

INRAN-SCAL: Italian National Food Consumption Survey

PNDS: National Demography and Health Survey

ENSANUT: Encuesta Nacional de Salud y Nutrición

ENSIN: Encuesta Nacional de Situación Nutricional

INCA 2: Individual and national study on food consumption

NDNS: British National Diet and Nutrition Survey

ORISCAV-LUX: Observation of Cardiovascular Risk Factors in Luxembourg

DNFCS: Dutch National Food Consumption Survey

ENASEM: Estudio Nacional de Salud y Envejecimiento en México

ENIDE: Encuesta Nacional de Ingesta Dietética

ENSO 2: Segunda Encuesta Nacional de Sobrepeso y Obesidad

NNPAS: National Nutrition and Physical Activity Survey

NEMONIT: German National Nutrition Monitoring

PNS: National Health Research

PDS: Research Social Dimension of Inequalities

SHIELD: Study to Help Improve Early Evaluation and Management of Risk Factors Leading to Diabetes

FFQ: Food Frequency Questionnaire

R24h: 24-hour dietary recall

WHO: World Health Organization

NCD: Noncommunicable diseases

DEXA: Dual-energy x-ray absorptiometry

References

- Acosta, K. (2013). Socioeconomic status and obesity in Colombia La obesidad y su concentración según nivel socioeconómico en Colombia. *Revista de Economía Del Rosario*, 16(2), 171–200.
- Acosta S.J. et al (2005). Estado nutricional de la población cubana adulta. *Rev Esp Nutr Comunitaria*, 11(1), 18–26.
- AHRQ. Agency for Healthcare Research and Quality. (n.d.). Newcastle-Ottawa Quality Assessment Form for Cohort Studies [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK115843/bin/appe-fm3.pdf>
- Alkerwi, A. et al (2015). Association between nutritional awareness and diet quality: Evidence from the observation of cardiovascular risk factors in Luxembourg (ORISCAV-LUX) study. *Nutrients*, 7(4), 2823–2838. <https://doi.org/10.3390/nu7042823>
- Alkerwi, A. et al. (2015a). *Cross-comparison of diet quality indices for predicting chronic disease risk: findings from the Observation of Cardiovascular Risk Factors in Luxembourg (ORISCAV-LUX) study*. 113(2), 259–269. <https://doi.org/10.1126/science.aaa1823>.
- Barquera, S. et al. (2009). Obesity and central adiposity in Mexican adults: results from the Mexican National Health and Nutrition Survey 2006. *Salud Publica de Mexico*, 51 Suppl 4, S595-603.
- Barquera, S. et al. (2010). Hypertension in Mexican adults: results from the National Health and Nutrition Survey 2006. *Salud Publica de Mexico*, 52 Suppl 1, S63-71. <https://doi.org/10.1590/S0036-36342010000700010>
- Barr, S. I., DiFrancesco, L., & Fulgoni, V. L. (2016). Association of breakfast consumption with body mass index and prevalence of overweight/obesity in a nationally-representative survey of Canadian adults. *Nutrition Journal*, 15(33), 2–9. <https://doi.org/10.1186/s12937-016-0151-3>
- Barros, M. B. de A. (2008). Inquéritos domiciliares de saúde: potencialidades e desafios. *Revista Brasileira de Epidemiologia*, 11(suppl 1), 6–19. <https://doi.org/10.1590/S1415-790X2008000500002>
- Bays, H. E., Chapman, R. H., & Grandy, S. (2007). The relationship of body mass index to diabetes mellitus, hypertension and dyslipidaemia: Comparison of data from two national surveys. *International Journal of Clinical Practice*, 61(5), 737–747. <https://doi.org/10.1111/j.1742-1241.2007.01336.x>
- Bell, L. K., Edwards, S., & Grieger, J. A. (2015). The relationship between dietary patterns and metabolic health in a representative sample of adult Australians. *Nutrients*, 7(8), 6491–6505. <https://doi.org/10.3390/nu7085295>
- Beltrán-de-Miguel, B., Estévez-Santiago, R., & Olmedilla-Alonso, B. (2015). Assessment of dietary vitamin A intake (retinol, α -carotene, β -carotene, β -cryptoxanthin) and its sources in the National Survey of Dietary Intake in Spain (2009–2010). *International Journal of Food Sciences and Nutrition*, 66(6), 706–712. <https://doi.org/10.3109/09637486.2015.1077787>
- Bezerra, I. N., & Sichieri, R. (2011). Household food diversity and nutritional status among adults in Brazil. *International Journal of Behavioral Nutrition and Physical Activity*, 8(22), 1–7. <https://doi.org/10.1186/1479-5868-8-22>
- Bjermo H. et al (2013). Lead, mercury, and cadmium in blood and their relation to diet among Swedish adults. *Food and Chemical Toxicology*, 57, 161–169.
- Cifelli, C. J., et al (2016). Increasing plant based foods or dairy foods differentially affects nutrient intakes: Dietary scenarios using NHANES 2007–2010. *Nutrients*, 8(7), 1–13. <https://doi.org/10.3390/nu8070422>
- Cook, Z. et al. (2005). Use of BMI in the assessment of undernutrition in older subjects: reflecting on practice. *The Proceedings of the Nutrition Society*, 64(3), 313–317. <https://doi.org/10.1079/PNS2005437>
- Cooper, R. et al. (2014). Body mass index from age 15 years onwards and muscle mass, strength, and quality in early old age: Findings from the MRC national survey of health and development. *Journals of Gerontology - Series A Biological Sciences and Medical Sciences*, 69(10), 1253–1259. <https://doi.org/10.1093/gerona/glu039>
- De Onis M, H. J. (1996). Anthropometric reference data for international use: recommendations from a World Health Organization Expert Committee. *Am. J. Clin. Nutr.*, 64, 650–658.
- Duarte, A. C. G. (2007). *Avaliação nutricional: aspectos clínicos e laboratoriais*. São Paulo: Atheneu.

- FAO. (2018). *Dietary assessment a resource guide to method selection and application in low resource settings*. <http://www.fao.org/3/i9940en/I9940EN.pdf>
- Ferreira, R. A. B., & Benicio, M. H. D. (2015). Obesidade em mulheres brasileiras : associação com paridade e nível socioeconômico. *Rev Panam Salud Publica*, 37(4/5), 337–342. <http://www.scielosp.org/pdf/tpsp/v37n4-5/v37n4-5a22.pdf>
- Fisberg, R. M., Marchioni, D. M. L., & Colucci, A. C. A. (2009). Avaliação do consumo alimentar e da ingestão de nutrientes na prática clínica. *Arquivos Brasileiros de Endocrinologia & Metabologia*, 53(5), 617–624. <https://doi.org/10.1590/S0004-27302009000500014>
- Gazan, R. et al (2016). Drinking Water Intake Is Associated with Higher Diet Quality among French Adults. *Nutrients*, 8(689), 2–23. <https://doi.org/10.3390/nu8110689>
- Gose, M. et al. (2016). Trends in food consumption and nutrient intake in Germany between 2006 and 2012: results of the German National Nutrition Monitoring (NEMONIT). *British Journal of Nutrition*, 115(08), 1498–1507. <https://doi.org/10.1017/S0007114516000544>
- Gray-Donald, K., Jacobs-Starkey, L., & Johnson-Down, L. (2000). Food habits of Canadians: Reduction in fat intake over a generation. *Canadian Journal of Public Health*, 91(5), 381–385.
- Guedes, F. A. G. (2007). *Avaliação nutricional de coletividades*. (4th ed.). Florianópolis: Editora da UFSC.
- Guendelman, S. et al (2013). A population-based comparison of weight and weight perceptions among overweight and obese Mexican and Mexican-American men TT - Comparación de base poblacional sobre el peso y percepciones sobre el peso entre hombres mexicanos y mexicano-americano. *Salud Publica Mex*, 55(supl.4), s451----s458.
- Hoffmann, K. et al (2002). Estimating the distribution of usual dietary intake by short-term measurements. *European Journal of Clinical Nutrition*, 56, S53–S62. <https://doi.org/10.1038/sj.ejcn.1601429>
- Nacional Cancer Institute (2015). Usual dietary intakes: The NCI method [Internet].
- Kant, A K; Graubard, B. (2015). 40-year trends in meal and snack eating behaviors of American adults. 115(5), 213–223. <https://doi.org/10.1007/978-1-62703-673-3>
- Kordas, K. et al (2012). Being Overweight or Obese Is Associated with Lower Prevalence of Anemia among Colombian Women of Reproductive Age 1 , 2. *The Journal of Nutrition*, (C), 175–181. <https://doi.org/10.3945/jn.112.167767.175>
- Kruizenga H.M. et al. (2003). Screening of nutritional status in The Netherlands. *Clinical Nutrition*, 22(2), 147–152. <https://doi.org/10.1054/clnu.2002.0611>
- Leclercq, C. et al. (2009). The Italian National Food Consumption Survey INRAN-SCAI 2005–06: main results in terms of food consumption. *Public Health Nutrition*, 12(12), 2504–2532. <https://doi.org/DOI: 10.1017/S1368980009005035>
- Lei, L. et al (2016). Dietary intake and food sources of added sugar in the Australian population. *British Journal of Nutrition*, 115, 868–877. <https://doi.org/10.1017/S0007114515005255>
- Lipschitz, D. (1994). Screening for nutritional status in the elderly. *Prim Care*, 21, 55–67.
- Malta, D. C., Célia, I., & Szwarcwald, L. (2017). Pesquisas de base populacional e o monitoramento das doenças crônicas não transmissíveis. *Inquéritos Apresentação Rev Saude Publica*, 51(1), 1S-4S. <https://doi.org/10.1590/S1518-8787.201705100supl1ap>
- Malta, D. C. et al (2008). Inquéritos Nacionais de Saúde: experiência acumulada e proposta para o inquérito de saúde brasileiro. *Revista Brasileira de Epidemiologia*, 11(supl 1), 159–167. <https://doi.org/10.1590/S1415-790X2008000500017>
- Marcenes, W. et al (2003). The relationship between dental status, food selection, nutrient intake, nutritional status, and body mass index in older people. *Cadernos de Saude Publica*, 19(3), 809–816. <https://doi.org/10.1590/S0102-311X2003000300013>
- Meier, M. et al (2010). Who eats healthily? A population-based study among young Swiss residents. *Public Health Nutrition*, 13(12), 2068–2075. <https://doi.org/10.1017/S1368980010000947>
- Melano-Carranza, E., Lasses Ojeda, L. A., & Ávila-Funes, J. A. (2008). Factores asociados con la hipertensión no tratada en los adultos mayores: resultados del estudio nacional sobre salud y envejecimiento en México, 2001^ies. *Rev. Panam. Salud Pública*, 23(5), 295–302.
- Meller, F.O et al. (2014). Associação entre circunferência da cintura e índice de massa corporal de mulheres brasileiras. *Ciência & Saúde Coletiva*, 19(1), 75–82. <https://doi.org/10.1590/1413-81232014191.2000>
- Moher, D. et al. (2015). Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews*, 4(1), 1–9. <https://doi.org/10.1186/2046-4053-4-1>
- Mohr, P. et al. (2007). Personal and lifestyle characteristics predictive of the consumption of fast foods in Australia. *Public Health Nutrition*, 10(12), 1456–1463. <https://doi.org/10.1017/S1368980007000109>
- Morimoto, J. M. et al. (2011). Within-person variance for adjusting nutrient distribution in epidemiological studies. *Revista de Saúde Pública*, 45(3), 621–625. <http://www.ncbi.nlm.nih.gov/pubmed/21552760>
- Nacif, M. (2007). *Avaliação antropométrica nos ciclos da vida: uma visão prática*. São Paulo: Editora Metha.

- Nissensohn, M. et al. (2017). Beverage consumption habits among the European population: Association with total water and energy intakes. *Nutrients*, 9(383), 1–13. <https://doi.org/10.3390/nu9040383>
- Nissensohn, M. et al. (2016). Beverage consumption habits and association with total water and energy intakes in the Spanish population: Findings of the ANIBES study. *Nutrients*, 8(232), 2–18. <https://doi.org/10.3390/nu8040232>
- Pavão, M. R. (2013). Autoavaliação do estado de saúde e a associação com fatores sociodemográficos, hábitos de vida e morbidade na população: um inquérito nacional. *Cad Saude Publica*, 29(4), 723–734. <https://doi.org/10.1590/S0102-311X2013000800010>
- Pereira RA, Duffey KJ, Sichieri R, P. B. (2014). Sources of excessive saturated fat, trans fat and sugar consumption in Brazil: an analysis of the first Brazilian nationwide individual dietary survey. *Public Health Nutr*, 17(1), 113–121. <https://doi.org/10.1017/S1368980012004892>
- Pisabarro, R. et al. (2009). Segunda Encuesta Nacional de Sobre peso y Obesidad (ENSO 2) adultos (18-65 años o más). *Revista Medica de Uruguay*, 25, 14–26.
- Ponce, X. et al (2013). Dietary quality indices vary with sociodemographic variables and anthropometric status among Mexican adults: a cross-sectional study. Results from the 2006 National Health and Nutrition Survey. *Public Health Nutrition*, 17(08), 1717–1728. <https://doi.org/10.1017/S1368980013002462>
- Pot, G K, et al. (2015). Trends in food consumption over 30 years: evidence from a British birth cohort. *European Journal of Clinical Nutrition*, 69(7), 817–823. <https://doi.org/10.1038/ejcn.2014.223>
- Pot, G. K, et al. (2014). Irregular consumption of energy intake in meals is associated with a higher cardio-metabolic risk in adults of a British birth cohort. *International Journal of Obesity*, 38(12), 1518–1524. <https://doi.org/10.1038/ijo.2014.51>
- Pot, G. K. et al (2014). Development of the Eating Choices Index (ECI): a four-item index to measure healthiness of diet. *Public Health Nutrition*, 17(12), 2660–2666. <https://doi.org/10.1017/S1368980013003352>
- Prynne, C. J. et al (2009). Meat consumption after disaggregation of meat dishes in a cohort of British adults in 1989 and 1999 in relation to diet quality. *European Journal of Clinical Nutrition*, 63(5), 660–666. <https://doi.org/10.1038/ejcn.2008.7>
- Rezende, F. et al (2007). Revisão crítica dos métodos disponíveis para avaliar a composição corporal em grandes estudos populacionais e clínicos. *ARCHIVOS LATINOAMERICANOS DE NUTRICION*, 57(4), 327–334.
- Ruiz, E. et al (2016). Macronutrient Distribution and Dietary Sources in the Spanish Population: Findings from the ANIBES Study. *Nutrients*, 7(6), 4739–4762. <https://doi.org/10.3390/nu7064739>
- Ruiz, E. et al (2015). Energy intake, profile, and dietary sources in the Spanish population: Findings of the ANIBES study. *Nutrients*, 7, 4739–4762. <https://doi.org/10.3390/nu7064739>
- Sette, S. et al (2018). The third Italian National Food Consumption Survey, INRAN-SCAI Part 1: Nutrient intakes in Italy. *Nutrition, Metabolism and Cardiovascular Diseases*, 21(12), 922–932. <https://doi.org/10.1016/j.numecd.2010.03.001>
- Shetty, P. S., & James, W. P. (1994). Body mass index. A measure of chronic energy deficiency in adults. *FAO Food and Nutrition Paper*, 56, 1–57.
- Shi, J., Ying, H., Du, J., & Shen, B. (2017). Serum Sclerostin Levels in Patients with Ankylosing Spondylitis and Rheumatoid Arthritis: A Systematic Review and Meta-Analysis. *BioMed Research International*, 2017, 1–7. <https://doi.org/10.1155/2017/9295313>
- Shim, J.-S., Oh, K., & Kim, H. C. (2014). Dietary assessment methods in epidemiologic studies. *Epidemiology and Health*, 36, 1–8. <https://doi.org/10.4178/epih/e2014009>
- Sichieri, R. et al. (2015). Major food sources contributing to energy intake – a nationwide survey of Brazilians aged 10 years and older. *British Journal of Nutrition*, 113(10), 1638–1642. <https://doi.org/10.1017/S0007114515001075>
- Silva, D. F. de O., Lyra, C. de O., & Lima, S. C. V. C. (2016). Padrões alimentares de adolescentes e associação com fatores de risco cardiovascular: uma revisão sistemática. *Ciência & Saúde Coletiva*, 21(4), 1181–1196. <https://doi.org/10.1590/1413-81232015214.08742015>
- Silveira, E. A., Kac, G., & Barbosa, L. S. (2009). Prevalência e fatores associados à obesidade em idosos residentes em Pelotas, Rio Grande do Sul, Brasil: classificação da obesidade segundo dois pontos de corte do índice de massa corporal. *Cad Saude Publica*, 25(7), 1569–1577. <https://doi.org/10.1590/S0102-311X2009000700015>
- Sluik, D., Engelen, A. I., & Feskens, E. J. (2014). Fructose consumption in the Netherlands: the Dutch national food consumption survey 2007–2010. *European Journal Of Clinical Nutrition*, 69, 475.
- Sluik, D. et al. (2014). Alcoholic beverage preference and diet in a representative Dutch population: the Dutch national food consumption survey 2007–2010. *European Journal Of Clinical Nutrition*, 68, 287.
- Sperandio, N., & Priore, S. E. (2017). Inquéritos antropométricos e alimentares na população brasileira: importante fonte de dados para o desenvolvimento de pesquisas. *Ciência & Saúde Coletiva*, 22(2), 499–508. <https://doi.org/10.1590/1413-81232017222.07292016>
- Szwarcwald, C. L. et al (2014). Pesquisa Nacional de Saúde no Brasil: concepção e metodologia de aplicação. *Ciência & Saúde Coletiva*, 19, 333–342. <https://doi.org/10.1590/1413-81232014192.14072012>
- Tande, D. L., Magel, R., & Strand, B. N. (2009). Healthy Eating Index and abdominal obesity. *Public Health Nutrition*, 13(02), 208–214. <https://doi.org/10.1017/S1368980009990723>

Tressou, J. et al (2016). Fatty acid dietary intake in the general French population: are the French Agency for Food, Environmental and Occupational Health & Safety (ANSES) national recommendations met? *British Journal of Nutrition*, 116(11), 1966–1973. <https://doi.org/10.1017/S000711451600413X>

Velasquez-Melendez, G. et al (2015). Prevalência de saúde cardiovascular ideal na população brasileira - Pesquisa Nacional de Saúde (2013). *Revista Brasileira de Epidemiologia*, 18(2013), 97–108. <https://doi.org/10.1590/1980-5497201500060009>

Vernay, M. et al. (2009). Association of socioeconomic status with overall overweight and central obesity in men and women: the French Nutrition and Health Survey 2006. *BMC Public Health*, 9, 215. <https://doi.org/10.1186/1471-2458-9-215>

Viacava, F. (2002). *Redalyc Informações em saúde : a importância dos inquéritos populacionais Health information : the relevance of health surveys*. 7(4), 607–621.

Whitton C. et al. (2011). National Diet and Nutrition Survey: fat and fatty acid intake from the first year of the rolling programme and comparison with previous surveys. In *British Journal of Nutrition* (Vol. 106). <https://doi.org/10.1017/S0007114511002911>

WHO. (2002). Active Ageing: A Policy Framework. *The Aging Male*, 5(1), 1–37. <https://doi.org/10.1080/713604647>

Willet, W. (2013). *Nutritional Epidemiology* (3rd ed.). New York: Oxford University Press.

World Health Organization. (2000). Obesity: Preventing and managing the global epidemic. Report of a WHO consultation. WHO Technical Report Series 894. *World Health Organization Technical Report Series*, 894, 1–253. <http://www.ncbi.nlm.nih.gov/pubmed/11234459>

World Health Organization. (2005). Envelhecimento Ativo: Uma política de saúde. *Organização Pan Americana de Saúde*, 1–62. [https://doi.org/10.1016/S0140-6736\(12\)61031-9](https://doi.org/10.1016/S0140-6736(12)61031-9)